

Letters

RESEARCH LETTER

PHYSICIAN WORK ENVIRONMENT AND WELL-BEING

Prevalence of Sexual Harassment in Academic Medicine

Sexual harassment is a form of gender discrimination that affects women and men in all areas of work.¹ According to the International Labour Organization (ILO), sexual harassment can occur in 1 or more of 3 forms: verbal, nonverbal, or physical.¹



Related article

Sexual harassment can lead to physical and psychological symptoms and diseases as well as work-related consequences.^{2,3} The prevalence of sexual harassment in medicine has been scantily investigated, and reports differ widely in the applied methodology.

Methods | All physicians working at a tertiary referral center in Berlin, Germany (n = 1862) were invited to participate in the survey for the current study between May 2015 and July 2015. This investigation was explicitly and solely designed to investigate sexual harassment. Institutional ethical approval (EA1/350/14, December 15, 2014) was obtained from Charité-Universitätsmedizin as well as from the employee representations/staff councils (clinical, academic and general), and all participants provided their written informed consent.

The survey instrument consisted of 36 items. It was administered online using the survey tool SoSci Survey (SoSci Survey GmbH). We investigated the following: (1) forms of misconduct experienced and whether these were considered harassing or threatening (note that not all misconduct was interpreted as harassment), (2) consequences experienced, (3) perpetrator profiles, (4) structural and organizational information, and (5) training and knowledge about sexual harassment, and (6) assumptions about its causes. Several questions about the type of misconduct or harassment experienced, the perpetrator profiles, and the consequences experienced were adapted from previous studies in German-speaking countries.^{4,5}

We computed descriptive statistics for sociodemographic and prevalence data. Prevalence and perpetrator profiles were compared bivariately between women and men using the Pearson χ^2 test. Multivariate regressions were calculated for structural factors using harassment patterns (any form, physical, nonphysical) as outcomes. We included only those participants who provided information on all investigated variables. All analyses were 2-sided, and $P < .05$ was considered significant. We performed all analyses using STATA software, version 13 (StataCorp LLC).

Results | A total of 790 physicians participated in the study (42% overall response rate), but 47 did not provide detailed infor-

Table 1. Experiences of Sexual Harassment Among Participating Physicians

Career Harassment and Misconduct Experiences ^a	Participants, No. (%)			P Value
	Total (n = 737)	Women (n = 448)	Men (n = 289)	
Any form	519 (70.04)	341 (76.12)	178 (61.59)	<.001
Degrading speech/obscene language ^b	456 (61.87)	297 (66.29)	159 (55.02)	.002
Letter, email, text message, jokes, pictures ^b	44 (5.97)	21 (4.69)	23 (7.96)	.07
Sexualized speech, sexual innuendo ^b	185 (25.1)	145 (32.37)	40 (13.84)	<.001
Whistling, staring ^b	96 (13.03)	89 (19.87)	7 (2.42)	<.001
Obscene gestures ^b	39 (5.29)	24 (5.36)	15 (5.19)	.92
Stories with sexual content ^b	109 (14.79)	79 (17.62)	30 (10.38)	.01
Sexual offers, unwanted invitations ^b	53 (7.19)	40 (8.93)	13 (4.5)	.02
Unwanted physical contact ^c	127 (17.23)	100 (22.32)	27 (9.34)	<.001
Groping, attempted kissing ^c	16 (2.17)	13 (2.9)	3 (1.04)	.09
Advantages for sexual favors	11 (1.49)	9 (2.01)	2 (0.69)	.15
Sexual attack ^c	4 (0.54)	2 (0.45)	2 (0.69)	.66
Other situations	15 (2.04)	11 (2.46)	4 (1.38)	.32
None of the above	218 (29.58)	107 (23.88)	111 (38.41)	<.001
Perceived the nonphysical misconduct as harassing (n = 492)	373 (75.8)	270 (83.3)	103 (61.3)	<.001
Perceived the nonphysical misconduct as threatening (n = 491)	59 (12)	49 (15.2)	10 (6)	.003
Perceived the physical misconduct as harassing (n = 132)	117 (88.6)	95 (93.1)	22 (73.3)	.003
Perceived the physical misconduct as threatening (n = 131)	36 (27.5)	31 (30.7)	5 (16.7)	.13

^a Multiple answers were accepted.

^b Summarized as nonphysical misconduct or harassment.

^c Summarized as physical misconduct or harassment.

Table 2. Perpetrator-Related and Structural Factors Correlated With Harassment Experiences

Characteristic	Harassed Individuals, No. (%)					
	Female (n = 333)			Male (n = 168)		
	Any Form	Nonphysical	Physical	Any Form	Nonphysical	Physical
Perpetrator Characteristics^a						
Mostly male	287 (86.2) ^b	272 (85.5) ^b	96 (95.1) ^b	63 (37.5) ^b	63 (38.4) ^b	4 (13.3) ^b
Mostly female	1 (0.3) ^b	1 (0.3) ^b	0	32 (19.1) ^b	28 (17.1) ^b	18 (60) ^b
Both male and female	45 (13.5) ^b	45 (14.1) ^b	5 (5) ^b	73 (43.5) ^b	73 (44.5) ^b	8 (26.7) ^b
Patients ^c	111 (32.6)	111 (34.2) ^d	39 (38.2)	45 (25.3)	43 (25.6) ^d	13 (43.3)
Relatives of patients ^c	37 (10.9)	36 (11.1)	14 (13.7)	14 (7.9)	14 (8.3)	3 (10)
Colleagues ^c	242 (71) ^d	237 (73.2) ^e	68 (66.7)	143 (80.3) ^d	141 (83.9) ^e	24 (80)
Superiors ^c	127 (37.2) ^b	120 (37) ^b	61 (59.8) ^b	32 (18) ^b	31 (18.5) ^b	3 (10) ^b
Structural Characteristics^f						
	Female Participants (n = 370)			Male Participants (n = 189)		
	Any Form	Nonphysical	Physical	Any Form	Nonphysical	Physical
Employment duration [≤5 years]	0.8 (0.5-1.3)	0.8 (0.5-1.2)	0.8 (0.5-1.2)	0.5 (0.2-0.9)	0.4 (0.2-0.8)	0.3 (0.2-0.9)
P value	.40	.30	.30	.02	.01	.05
Immediate team size [≥11 people]	1.4 (0.8-2.3)	1.4 (0.8-2.3)	1.2 (0.8-2.1)	1.0 (0.5-2.0)	1.0 (0.6-2.0)	1.8 (0.6-5.1)
P value	.20	.20	.40	.90	.90	.30
Institute/department size [≥41 people]	0.9 (0.5-1.4)	0.9 (0.5-1.4)	0.9 (0.5-1.4)	0.7 (0.4-1.4)	0.9 (0.5-1.8)	0.7 (0.2-1.9)
P value	.60	.60	.60	.40	.80	.50
Sex distribution team [mostly men]	1.5 (0.8-2.9)	1.8 (0.9-3.6)	0.7 (0.4-1.4)	1.6 (0.7-4.0)	1.7 (0.7-4.3)	2.3 (0.7-7.2)
P value	.20	.07	.40	.30	.20	.20
Sex of team leader [man]	1.4 (0.9-2.4)	1.6 (0.9-2.6)	0.9 (0.5-1.5)	1.0 (0.4-2.2)	0.9 (0.4-1.9)	1.4 (0.4-5.1)
P value	.20	.08	.60	>.99	.70	.60
Sex of department leader [man]	0.7 (0.4-1.3)	1.0 (0.6-1.7)	0.7 (0.4-1.1)	1.5 (0.7-3.2)	1.4 (0.7-3.0)	2.2 (0.6-8.6)
P value	.30	.90	.10	.30	.30	.20
Hierarchy [strong hierarchy]	1.3 (0.8-2.0)	1.3 (0.8-2.0)	1.9 (1.1-3.1)	2.3 (1.2-4.4)	2 (1-3.8)	2.6 (0.9-7.3)
P value	.60	.30	.01	.01	.04	.06
Knowledge of available recourse inside the hospital [no knowledge]	0.7 (0.4-1.3)	0.7 (0.4-1.3)	0.8 (0.4-1.4)	1.3 (0.6-2.9)	1.5 (0.7-3.2)	1.1 (0.3-3.5)
P value	.30	.30	.40	.50	.30	.90
Knowledge of available recourse outside the hospital [no knowledge]	1.4 (0.8-2.5)	1.2 (0.7-2.2)	1.4 (0.8-2.6)	0.7 (0.3-1.7)	0.9 (0.4-2.1)	0.8 (0.2-3.0)
P value	.30	.50	.20	.50	.80	.80
Ever had formal anti-harassment training [never had training]	1.0 (0.5-2.2)	1.0 (0.5-2.2)	0.7 (0.3-1.5)	1.6 (0.6-4.3)	1.5 (0.5-4)	2.5 (0.4-14)
P value	.90	.90	.40	.40	.40	.30

Abbreviation: OR, odds ratio.

^a Perpetrator characteristics are expressed as absolute values (percentages) and have been analyzed as female-male bivariate comparisons using the Pearson χ^2 test.

^b $P < .001$.

^c Multiple answers accepted.

^d $P < .05$.

^e $P < .01$.

^f Structural factors are expressed as OR (95% CI) and have been analyzed segregated by sex using multivariate models adjusted for all presented covariables. Reference outcomes are reported in brackets.

mation; 2 did not provide information on sex; and 4 (1%) self-reported as transgender, intersex, or nonbinary. The low number of individuals self-reporting other than distinct male or female gender prevented statistical testing of this category, but it should be noted that all 4 of these participants (100%) reported experiencing some form of harassment. Of the 737 participants included in the full analysis, 60% were women (n = 448), and 39% were men (n = 289).

All of the absolute numbers for the following reports can be found in **Table 1**. Among all male and female participants, 70% reported some form of misconduct while performing their

work. The most common form self-reported as harassment was verbal harassment (including degrading speech [62%] and sexualized speech [25%]). Nonphysical misconduct was perceived as harassing by 76% of the individuals, more frequently by women than by men (83% vs 61%; $P < .001$). Physical misconduct was perceived as harassing by 89% of those reporting such misconduct and as threatening by 28%, with no significant sex differences (Table 1).

Women reported the perpetrators of harassment to be almost exclusively male, both for nonphysical harassment (85% of perpetrators against women compared with 38% of pepe-

trators against men; $P < .001$) (Table 2) and for physical harassment (95% of perpetrators against women compared with 13% of those against men; $P < .001$) (Table 2). Colleagues were reported as the main perpetrators at similar rates by men and women, while women reported superiors to be the perpetrators more frequently (37% vs 18%; $P < .001$) (Table 2).

Strong departmental or divisional hierarchy appeared as the only structural factor significantly associated with harassment in both male and female victims (all supporting data reported in Table 2).

Discussion | Sexual harassment frequently affects female and male physicians during their careers. In the present study, both groups reported verbal harassment as the most frequent form of misconduct. While perpetrator patterns differed between male and female victims, strong institutional hierarchies were associated with sexual harassment in both sexes, highlighting the importance of organizational culture.⁶ These results support the need for cultural change in the form of structural and widespread action to truly reduce the high incidence of sexual harassment in academic medicine.

Sabine Jenner, BA, MA

Pia Djermester, BA

Judith Prügl, MD

Christine Kurmeyer, PhD

Sabine Oertelt-Prigione, MD, MScPH, PhD

Author Affiliations: Equal Opportunity Office, Charité-Universitätsmedizin, Berlin, Germany (Jenner, Djermester, Prügl, Kurmeyer); Department of Primary and Transmural Care, Radboudumc, Nijmegen, the Netherlands (Oertelt-Prigione); Institute of Legal Medicine, Charité-Universitätsmedizin, Berlin, Germany (Oertelt-Prigione).

Accepted for Publication: August 4, 2018.

Corresponding Author: Sabine Oertelt-Prigione, MD, PhD, MSc, Chair of Gender in Primary and Transmural Care, Department of Primary and Community Care, Radboud Institute of Health Sciences, Radboud University Medical Center, Geert Grooteplein 21 (route 117), 6500HB Nijmegen, The Netherlands (sabine.oertelt-prigione@radboudumc.nl).

Published Online: October 3, 2018. doi:10.1001/jamainternmed.2018.4859

Author Contributions: Dr Oertelt-Prigione had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Jenner, Kurmeyer, Oertelt-Prigione.

Acquisition, analysis, or interpretation of data: Jenner, Djermester, Prügl, Oertelt-Prigione.

Drafting of the manuscript: Oertelt-Prigione.

Critical revision of the manuscript for important intellectual content: Jenner, Djermester, Prügl, Kurmeyer.

Statistical analysis: Oertelt-Prigione.

Obtained funding: Kurmeyer, Oertelt-Prigione.

Administrative, technical, or material support: Djermester, Prügl, Kurmeyer.

Study supervision: Jenner, Oertelt-Prigione.

Conflict of Interest Disclosures: None reported.

Funding/Support: This work has been supported by the Charité Foundation, Berlin, Germany; the Equal Opportunity Program of the City of Berlin, Berlin, Germany; and the Hans Böckler Stiftung, Duesseldorf, Germany.

Role of the Funder/Sponsor: The funding organizations had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Meeting Presentation: This article was presented at the North American Menopause Society Annual Meeting; October 3-6, 2018; San Diego, California.

Additional Contributions: We thank all the participants, who have generously provided us with their time and have shared their sometimes difficult experiences.

1. International Labour Organization. Code of Conduct on Sexual Harassment in the Workplace Brochure; 2015. http://www.ilo.org/global/docs/WCMS_371182/lang-en/index.htm. Accessed September 13, 2018.

2. Bowling NA, Beehr TA. Workplace harassment from the victim's perspective: a theoretical model and meta-analysis. *J Appl Psychol*. 2006;91(5):998-1012. doi:10.1037/0021-9010.91.5.998

3. Willness CR, Steel P, Lee K. A meta-analysis of the antecedents and consequences of workplace sexual harassment. *Personnel Psychol*. 2007;60(1):127-162. doi:10.1111/j.1744-6570.2007.00067.x

4. Mueller U, Schroettle M. Lebenssituation, Sicherheit und Gesundheit von Frauen in Deutschland; 2004. <https://www.bmfsfj.de/blob/jump/84328/langfassung-studie-frauen-teil-eins-data.pdf>. Accessed September 13, 2018.

5. Krings F, Schär Moser M, Mouton A. *Sexuelle Belästigung am Arbeitsplatz: Wer belästigt wen, wie und warum? Besseres Verständnis heisst wirksamere Prävention*. Lausanne, Switzerland: Université de Lausanne; 2013.

6. Frank E, Brogan D, Schiffman M. Prevalence and correlates of harassment among US women physicians. *Arch Intern Med*. 1998;158(4):352-358. doi:10.1001/archinte.158.4.352